

The disinformation problem for black holes

Sean A. Hayward

Department of Physics, National Central University, Jhongli, Taoyuan 320, Taiwan

No escape?

1966: “*event horizon*... is the boundary of the region from which particles or photons can escape to infinity... a *black hole* is a region... from which particles or photons cannot escape”

1976: “Because part of the information about the state of the system is lost down the hole, the final situation is represented by a density matrix rather than a pure quantum state”

1997: “Whereas Stephen Hawking and Kip Thorne firmly believe that information swallowed by a black hole is forever hidden from the outside universe, and can never be revealed even as the black hole evaporates and completely disappears...”

2004: “Thus the total path integral is unitary and information is not lost in the formation and evaporation of black holes. The way the information gets out seems to be that a true event horizon never forms, just an apparent horizon” [Hawking quotes]

How should black holes be defined? Event and apparent horizons are too indirectly defined.

Assume spherical symmetry for simplicity (everything generalizes).

Area A , null coordinates $x^\pm$: $g^{-1}(dx^\pm, dx^\pm) = 0$, future-pointing, unique up to $x^\pm \mapsto \tilde{x}^\pm(x^\pm)$.


Null expansions $\theta_\pm = \partial_\pm A/A$, $\partial_\pm = \partial/\partial x^\pm$.

Normally, outgoing light rays diverge, $\theta_+ > 0$,

ingoing light rays converge, $\theta_- < 0$;

outgoing wavefront expands,

ingoing wavefront contracts:



But if there is a massive source inside the surface,


its gravitational field has an attractive or converging effect.

Close enough to a massive enough source,



the outgoing light rays may also be converging, $\theta_+ < 0$; a **trapped surface**:

Everything inside is trapped within a shrinking area.

Nothing can escape, not even light.



In between, there will be a **marginal surface** (marginally trapped surface), $\theta_+ = 0$, where the outgoing light rays are instantaneously parallel:



This is a black hole: its surface is located by the marginal surface, where outgoing light rays are instantaneously parallel, $\theta_+ = 0$, ingoing light rays are converging, $\theta_- < 0$, and outgoing light rays are diverging just outside and converging just inside, $\partial_- \theta_+ < 0$. As time develops, the marginal surfaces generate a hypersurface in space-time.

Gravity traps

$$\text{A surface is } \begin{cases} \text{untrapped} \\ \text{marginal} \\ \text{trapped} \end{cases} \text{ if } \theta_+ \theta_- \begin{cases} < 0 \\ = 0 \\ > 0 \end{cases},$$


$$\begin{cases} \text{future} \\ \text{past} \end{cases} \text{ trapped if } \theta_{\pm} \begin{cases} < 0 \\ > 0 \end{cases}, \begin{cases} \text{future} \\ \text{past} \end{cases} \text{ marginal if } \theta_- \begin{cases} < 0 \\ > 0 \end{cases} \text{ (for } \theta_+ = 0 \text{)}.$$

Trapping horizon: a hypersurface foliated by marginal surfaces,

$$\begin{cases} \text{outer} \\ \text{degenerate} \\ \text{inner} \end{cases} \text{ if } \partial_- \theta_+ \begin{cases} < 0 \\ = 0 \\ > 0 \end{cases} \text{ (for } \theta_+ = 0 \text{)}.$$

A $\begin{cases} \text{future} \\ \text{past} \end{cases}$ outer trapping horizon can be taken as the local definition of a $\begin{cases} \text{black} \\ \text{white} \end{cases}$ hole.

e.g. Schwarzschild and Reissner-Nordström black holes:



Signature law:

$$\text{NEC (null energy condition)} \Rightarrow \begin{cases} \text{outer} \\ \text{inner} \end{cases} \text{ trapping horizons are } \begin{cases} \text{spatial or null} \\ \text{temporal or null} \end{cases},$$

and null if and only if the ingoing energy density vanishes.

Proof. Take the case $\theta_+ \cong 0$ (evaluation on the horizon),

write the horizon-generating vector $\xi = \xi^+ \partial_+ + \xi^- \partial_-$,


then $0 \cong \xi \cdot d\theta_+ = \xi^+ \partial_+ \theta_+ + \xi^- \partial_- \theta_+$ relates three signs:

$$\frac{\xi^-}{\xi^+} \begin{cases} < 0 \\ = 0 \\ > 0 \end{cases} \text{ for a } \begin{cases} \text{spatial} \\ \text{null} \\ \text{temporal} \end{cases} \text{ trapping horizon;}$$


$$\partial_- \theta_+ \begin{cases} < 0 \\ > 0 \end{cases} \text{ for an } \begin{cases} \text{outer} \\ \text{inner} \end{cases} \text{ trapping horizon;}$$

$$\partial_+ \theta_+ \cong -8\pi G T_{++} \text{ (null focusing equation, from Einstein equation),}$$

$$T_{\pm\pm} = T(\partial_{\pm}, \partial_{\pm}) \text{ (energy tensor), NEC} \Rightarrow T_{\pm\pm} \geq 0.$$



$\Rightarrow$ **Black hole horizons are one-way traversable;**
one can fall into a black hole but not escape:



Area law: NEC $\Rightarrow$

$$\left\{ \begin{array}{l} \text{future outer or past inner} \\ \text{past outer or future inner} \end{array} \right\} \text{trapping horizons have } \left\{ \begin{array}{l} \text{non-decreasing} \\ \text{non-increasing} \end{array} \right\} \text{area, } \left\{ \begin{array}{l} A' \geq 0 \\ A' \leq 0 \end{array} \right\},$$

instantaneously constant ($A' = 0$) if and only if the horizon is null.

Proof. $A' = \xi \cdot dA = A(\xi^+ \theta_+ + \xi^- \theta_-) \cong A \xi^- \theta_-$. Fixing orientation $\xi^+ > 0$,

$$\xi^- \left\{ \begin{array}{l} \leq 0 \text{ for outer horizons} \\ \geq 0 \text{ for inner horizons} \\ = 0 \text{ for null horizons} \end{array} \right\} \text{ as above; } \theta_- \left\{ \begin{array}{l} < 0 \\ > 0 \end{array} \right\} \text{ for } \left\{ \begin{array}{l} \text{future} \\ \text{past} \end{array} \right\} \text{ horizons.}$$

$\Rightarrow$ Black holes grow if they absorb any matter, and otherwise remain the same size.

(Actually general results, not restricted to spherical symmetry).


Evaporating black holes: Hawking radiation is based on pair production of positive-energy outgoing radiation, $T_{--} > 0$, and negative-energy ingoing radiation, $T_{++} < 0$.
 $\Rightarrow$ area and signature properties reverse:

$$\text{ingoing energy density negative} \Rightarrow \left\{ \begin{array}{l} \text{outer} \\ \text{inner} \end{array} \right\} \text{trapping horizons are } \left\{ \begin{array}{l} \text{temporal} \\ \text{spatial} \end{array} \right\},$$

$$\left\{ \begin{array}{l} \text{future outer or past inner} \\ \text{future inner or past outer} \end{array} \right\} \text{trapping horizons have } \left\{ \begin{array}{l} \text{decreasing} \\ \text{increasing} \end{array} \right\} \text{area, } \left\{ \begin{array}{l} A' < 0 \\ A' > 0 \end{array} \right\}.$$

The black-hole horizon is shrinking and two-way traversable:

Matter and information clearly will escape.




Losing it?

No reason to expect a purely thermal spectrum for an evaporating black hole;

Hawking's thermal spectrum holds only for a stationary black hole,

ignoring the back-reaction of the radiation on the black hole, which is not evaporating.

Endpoint of evaporation? Usual picture:



PO: $T_{++} > 0$, outer $\Rightarrow$ spatial, growing.

ON: $T_{++} < 0$, outer $\Rightarrow$ temporal, shrinking.

The horizon shrinks to zero size, intersecting the singularity.

Some of the matter (and information) escapes, some is lost in the singularity.

However, this is a semi-classical approximation, valid only away from the singularity.

In full quantum gravity, is the singularity resolved?

The great escape

Assume that the centre never becomes singular.

Regular centre $\Rightarrow \exists$ untrapped neighbourhood

$\Rightarrow \exists$ inner (future) trapping horizon.

Inner horizon and centre approach, become almost null,
matter becomes almost pure outgoing radiation:

($\exists$ exact Einstein-Klein-Gordon solution).

During black-hole formation:

PO: $T_{++} > 0$, outer $\Rightarrow$ spatial, growing.

PI: $T_{++} > 0$, inner $\Rightarrow$ temporal, shrinking.

During black-hole evaporation:

ON: $T_{++} < 0$, outer $\Rightarrow$ temporal, shrinking.


IN: $T_{++} < 0$, inner $\Rightarrow$ spatial, growing.

The outer and inner sections eventually meet,
marking the endpoint of evaporation.

Metric $C^2 \Rightarrow$ inner and outer sections join smoothly.

$\Rightarrow \exists$ single trapping horizon of $S^2 \times S^1$ topology,
enclosing a compact region of trapped surfaces.

The black hole has evaporated completely,
with all its contents re-emerging, albeit mangled.



No information loss; no event horizon. Obtained by (at least) three independent methods:

- Euclidean quantum gravity [Hawking, unpublished];
- loop quantum gravity [Ashtekar & Bojowald, unpublished];
- this qualitative classical analysis with minimal assumptions about “correct” quantum gravity,
that a singularity never forms
and that the ingoing radiation has negative energy density during the entire evaporation phase.

Orthodox paradox

- Logically there are no true paradoxes, just misunderstandings, usually simple but fundamental.

The *disinformation problem* for black holes: defining them by event horizons as regions of no escape.

- Understanding black holes locally by trapping horizons, the supposed paradox disappears.